

# Comparative Survey of Geometric Morphometrics and Traditional Methods in Male and Female of Anjak, *Schizocypris altidorsalis* (Bianco and Banarescu, 1982) from Sistan Basin, Iran

Zohreh Ganjali

Department of Biology, Faculty of Basic Science, University of Zabol, Zabol, Iran

Corresponding author's e-mail: zohreganjali@yahoo.com

## Article Information

Received: 10 August 2024  
Revised: 26 September 2024  
Accepted: 28 September 2024  
Published online: 14 February 2025

## Keywords

*Schizocypris altidorsalis*  
Phenotype plasticity  
Procrustes analysis  
Hotelling t-test  
Principal component analysis

## Abstract

Morphological variation between male and female populations of Anjak, *Schizocypris altidorsalis* species in Sistan basin of Iran was studied using landmark-based geometric morphometric and Traditional morphometric. A total of 27 specimens (16 males and 13 females) were caught from one basin. A total number of 16 landmark-points were defined and digitized on 2D pictures, and 29 morphometric characters were measured. To eliminate non-shape data, including size, direction, and position, the extracted data were analyzed by the generalized procrustes analysis. The data of TM were analyzed by Hotelling t-test, and data of GM were analyzed and compared using multivariate principal component analysis (PCA) which showed male and female populations overlapped. Deformation shape grid and Wireframe based on the consensus body shape, very little separation the two populations based on the head size, body depth, and caudal peduncle length characteristics. Based on TM analysis, these characters could not differentiate between the female and male, except for two characters, length of pectoral fin and maximum body depth ( $p < 0.00$ ). Comparative morphological studies revealed deformation of the grid and wireframe in more details. In the deformation grid, there are differences between the two populations due to folds and twists. Wireframe shows differences better in certain places, such as the head, caudal peduncle, and middle area of the body.

© 2025 University of Zabol. All rights reserved.

## 1. Introduction

Iran is located in the southern Palearctic biogeographic region bordering the Oriental and Ethiopian regions and is considered as exchange region for freshwater fishes [5], with 297 species in 109 genera and 30 families [9]. *Schizocypris altidorsalis*, belonging to the Cyprinidae family, is one of the economically valuable fishes. It is a

rare native species of the genus *Schizocypris*. It belongs to the subfamily Schizothoracinae, which is found in the southeastern regions of Iran, Central Asia and China, India, Pakistan, and Afghanistan [1]. The Sistan area is located in the southeastern of Iran.

Anjak (*Schizocypris altidorsalis*), is a native benthopelagic and commercially important fish species in Hamun Lake and Hirmand River system in Sistan basin, Iran [11]. Together with snow trout (*Schizothorax zarudnyi*), Anjak make up the most favorite source of animal protein for Sistan residents [11]. As an endemic species in the Sistan region of Iran, *Schizocypris altidorsalis* Bianco and Banarescu, 1982 (Cyprinidae: Schizothoracinae) resides only in the Hamoun Lake, Helmand River, and its tributaries and presumably adjacent Afghanistan [4]. The genus *Schizocypris* of mediumsized snow trouts contains only two species that were found in Pakistan, Afghanistan, and Iran. *S. altidorsalis* is a benthopelagic species that has been reported from pools in dry river beds and still, reedy channels. The fish enter the Hamoun Lake from the upstream parts of the rivers, and return to more permanent rivers when water levels fall [4,12].


For the first time, in this paper, we simultaneously performed traditional morphometric and geometric morphometrics measurements for *S.altidorsalis*. The study of the body shape is a routine method for understanding many aspects of fish biology, such as resource management, evolution, behavior, and ecology [25,15]. For instance, geometric morphometric methods revealed some phenotypic plasticity evidences related to dam construction in *Capoeta gracilis* [13]. Intraspecific morphological variations in fishes are related to the adaptations of organisms to the habitat environmental conditions [2,6]. Hence, morphological studies are among the tools used to identify different fish populations and better understanding of habitat-associated morphological divergence [21]. Geometric morphometrics (GM), a quantitative approach to the analysis of shape, is widely applied to compare and determine shape variations of biological structures [27].

Some authors have implemented the geometric morphometric method (GM) [7, 19, 20, 29, 28, 17, 18, 10]. It has been suggested that GM is more effective in detecting morphological disparities [14, 23, 10]. Additionally, it has been shown that both methods can lead to similar [17] or different results [27, 10]. In the paper by Ganjali et al. 2021, the geometric morphometric method (GM) than the traditional method showed more differences and was more accurate.

## 2. Materials and Methods

### 2.1 Sample collection and preparation

For this study, 29 fish samples (16 males and 13 females) representing Anjak were obtained from Chahnimeh reservoirs, Sistan, Iran (30°50'21.7500"N, 61°41'39.8004"E) by fishing (Figure 1). The specimens were preserved in 4% buffered formaldehyde after anesthesia in 1% clove oil solution.



**Figure 1.** Study area from Chahnimeh reservoirs in Sistan Basin

## 2.2 Traditional morphometric (TM)

29 morphometric characters were measured with a digital caliper (0.05mm accuracy). The elimination of the size effects from the data set was done based on Elliott et al. 1995 using the formula of  $M_{adj} = M (L_s/L_o)^b$ , where  $M_{adj}$  is the size adjusted measurement,  $M$  is the original measurement,  $L_s$  is overall mean of standard length for all fish from all samples in each analysis,  $L_o$  is the total length of fish parameter, and  $b$  was estimated for each character from the observed data as the slope of the regression of  $\log M$  on  $\log L_o$ , using all fish in all groups. Morphometric characters were analyzed for normality distribution using the Kolmogorov–Smirnov test and T- test in SPSS software version 22.

### 2.2.1 Traditional morphometric characters


Total Length, Fork Length, Standard Length, Head Length, Head depth, Head Width, Pre orbital distance, post orbital distance, Inter orbital distance, Eye diameter, Predorsal distance, Post-dorsal distance, Length of dorsal fin, Depth of dorsal fin, Length of Anal fin, Depth of Anal fin, Length of pectoral fin, Depth of pectoral fin, Length of ventral fin, Depth of ventral fin, Pre anal distance, Post anal distance, Distance between pectoral and ventral fin, Distance between Ventral and Anal fin, Length of Caudal fin, Depth of Caudal fin, Minimum body depth, Maximum body depth, Length of caudal peduncle, Mouth width.

## 2.3 Geometric morphometric (GM)

The purpose of geometric morphometry is to study the process of variability and body shape patterns. Geometric morphometrics is suitable for comparing populations because the number of traits we consider is more than the traditional method. The left sides of fish specimens were photographed using a copy stand equipped with a digital camera (Canon G7). To extract data on the body shape, 16 homologous landmark points were defined and digitized using tpsDig2 software (version 2.31) (Figure 2) on 2D pictures.

### 2.3.1 Geometric morphometric characters

The following Geometric morphometric characters were measured (Figure 2): (1) anterior-most point of the snout tip on the upper jaw, (2) the beginning of the eye socket, (3) center of the eye, (4) the end of the eye socket, (5) the dorsal surface of the head is perpendicular to the center of the eye, (6) the end point of the gill operculum, (7) the origin point of the dorsal-fin base, (8) the insertion point of the dorsal-fin base, (9) the end of the caudal peduncle or the beginning of the procrand radius, (10) the middle part of the most end of the caudal peduncle, (11) the lower part of the most recessed of the caudal peduncle, (12) the origin point of the anal-fin base, (13) the insertion point of the anal- fin base, (14) most anterior point of the pectoral fin, (15) the most ventral point of the gill slit and (16) ventral surface of the head perpendicular to the center of the eye.



**Figure 2.** Defined landmark points to extract the body shape data of *S. altidorsalis*

Then, to remove non-shape variations (location, orientation and scale) from the landmark configurations, they were superimposed by GPA (General Procrustes Analysis) [25]. Finally, the principle component analysis (PCA) was also conducted.

### 3. Results and Discussion

#### 3.1 Traditional morphometric analysis


Evaluate results and comparison of Traditional morphometric 29 specimens of Anjak (*Schizocypris altidorsalis*) shown in Table 1. In this study, number 13 female and 16 male numbers. In length of pectoral fin and maximum body depth  $p < 0.00$ .

**Table1.** Statistical mean and results of T-test measuring traits of male and female Anjak, *S. altidorsalis* (Bianco and Banarescu, 1982) from Sistan Basin, Iran


Characters (mm)	Male(n=16)	Female(n=13)	P value
<b>SL</b>	125.53	125.53	-
<b>HL</b>	29.26±1.69	28.45±1.10	0.147
<b>HD</b>	14.93±2.74	16.89±7.52	0.342
<b>Hwidth</b>	12.55±0.81	12.87±0.53	0.232
<b>Pre orbit Dis</b>	10.19±0.98	9.93±0.99	0.485
<b>IntOD</b>	9.85±0.52	9.94±0.48	0.636
<b>Post Orbit D</b>	14.57±1.22	13.74±1.04	0.636
<b>EyeDiam</b>	5.32±0.54	5.44±0.68	0.582
<b>Pre DD</b>	63.92±1.49	64.20±5.67	0.853
<b>PosDD</b>	76.52±5.21	77.75±3.38	0.47
<b>LDfin</b>	16.81±1.47	17.58±2.13	0.263
<b>DDfin</b>	30.05±1.36	30.02±2.46	0.963
<b>LAfin</b>	10.47±2.28	9.83±1.98	0.435
<b>DAfin</b>	24.05±3.14	24.05±2.84	0.998
<b>LPecfin</b>	19.51±1.87	20.86±0.90	0.026*
<b>DPecfin</b>	5.01±0.70	4.76±0.47	0.284
<b>LVentfin</b>	16.69±1.70	16.30±1.57	0.528
<b>Dventfin</b>	4.66±0.38	4.77±0.64	0.597
<b>PreAD</b>	97.06±2.43	97.09±2.86	0.969
<b>PosAD</b>	60.86±4.33	60.43±2.56	0.754
<b>Dis P&amp;V</b>	39.49±1.85	39.73±2.71	0.781
<b>Dis V&amp;A</b>	32.69±1.63	32.69±2.12	0.992
<b>LcaudF</b>	29.62±3.89	30.27±1.97	0.587
<b>MaxBD</b>	26.46±2.67	28.85±2.08	0.014*
<b>MinBD</b>	10.22±0.52	10.36±0.95	0.6
<b>LCaudPed</b>	15.35±2.07	15.36±1.55	0.989
<b>MWidth</b>	8.07±0.69	7.92±0.90	0.613

#### 3.2 Geometric morphometric analysis


Based on the results, the first four components of PCA being above the Jolliffe cutoff line (PC1= 40.9, PC2=23.81, PC3=9.66, PC4=8.63) Figure 3, were selected as the main factors separating female and male populations. As presented in Figure 4, the populations are grouped along the first two axes and the studied populations presented no in size but also overlapped. In deformation grid (Figure 5) and wire frame (Figure 6) changes are seen.




**Figure 3.** The screen plot diagram of PCA and the Jolliffe cutoff point (red line), showing the main (4) significant components Jolliffe line



**Figure 4.** Principal component analysis diagram of male and female Anjak, *S. altidorsalis* in the Sistan Basin



**Figure 5.** Comparison of the average body shape of male and female Anjak, *S. altidorsalis* in Sistan Basin using the deformation Grid network.



**Figure 6.** Comparison of the average body shape of male and female Anjak, *S. altidorsalis* in Sistan Basin using the deformation Wireframe network (Female: Red, Male: Blue)

Occurrence of prolonged droughts and Sistan region diminished spawning fish and uncontrolled fishing in this area Anjak the fear that goes with this species of value and that regional endemic economy fish disposable component of is considered at risk should be destroyed. This paper performed traditional morphometric and 2D landmark based geometric methodologies were used for the discrimination of one population of Anjak, *S. altidorsalis* collected from Sistan basin of Iran.

### 3.3 Traditional morphometrics

Based on TM analysis, these characters could not differentiate between female and male populations. Table 1 shows that the results were not significantly different  $p$  Value  $> 0$  except two characters (length pectoral fin=0.026 and maximum body depth=0.014). It has been suggested that the morphological characteristics of fish are determined by an interaction between genetic and environmental factors [25-15].

### 3.4 Geometric morphometrics

Based on the results (deformation grid and wireframe) of male and female populations, the head, caudal peduncle, depth of body and tail (Figures 5 & 6) were more affected parts in terms of morphological changes, indicating that this species uses these areas to move and adapt in different environments. The PCA analysis along the first two axes (PC1 and PC2) of geometric-based data for the *S. altidorsalis* population showed that (Figure 4) overlapped. Therefore, due to the overlap, there is no difference based on this test (PCA) between the two populations of Anjak males and females.

## 4. Conclusion

Comparative morphological studies (TM & GM) revealed deformation grid and wireframe with more details. In the deformation grid, there are differences between the two populations due to folds and twists. Wireframe shows differences better in certain places, such as the head, caudal peduncle, and middle area of the body. In general, the geometric results are more accurate and better than Traditional morphometric. I hope that the drought problem will be solved in the future so that we can study more populations for phenotypic adaptation.

## Conflicts of Interest

The author declares no conflict of interest associated with this manuscript.

## Acknowledgments

I would like to thank to the people who helped me for fish collection and A. Mouludi-Saleh for helping with analysis.

## References

1. Abbaspour R, Rahbar M, Mesgaran Karimi J. Comparative Survey of Morphometric-meristic Male and Female Anjak Fish (*Schizocypris brucei*, Annandale and Hora, 1920) of Hamoun Wetland in South East Iran. *Middle East J. Sci. Res.*, 2013, 14(5):620-623.
2. Bentzen P, McPhail JD. Ecology and evolution of sympatric sticklebacks (*Gasterosteus*)-specialization for alternative trophic niches in the Enos lake species pair. *Can. J. Zool.*, 1984, 62(11):2280-2286.
3. Coad BW. Zoogeography of the fishes of the Tigris-Euphrates basin. *Zool. Middle East*, 1996 13(1):51-70.

4. Coad BW. *Freshwater fishes of Iran*. 2013. www.briancoad.com
5. Coad BW. *Freshwater fishes of Iran*. 2019. www.briancoad.com
6. Cresko WA, Baker JA. Two morphotypes of lacustrine threespine stickleback, *Gasterosteus aculeatus*, in Benka Lake, Alaska. *Environ. Biol. Fishes*, 1996, 45(4):343-350.
7. Eagderi S, Esmailzadegan E, Madah A. Body shape variation in riffle minnows (*Alburnoides eichwaldii* De Filippii, 1863) populations of Caspian Sea basin. *Taxonomy and Biosystematics*, 2013, 5(14):1-8.
8. Elliott NG, Haskard K, Koslow JA. Morphometric analysis of orange roughy (*Hoplostethus atlanticus*) off the continental slope of southern Australia. *J. Fish Biol.*, 1995, 46:202e220.
9. Esmaili HR, Sayyadzadeh G, Eagderi S, Abbasi K. Checklist of freshwater fishes of Iran. *FishTaxa*, 2018, 3(3):1-95.
10. Ganjali Z, Esmaili HR, Eagderi S, Gholamhosseini A. Can geometric morphometrics and traditional methods work for evaluation of body shape plasticity of exotic fishes? A case study of topmouth gudgeon, *Pseudorasbora parva* (Teleostei: Gobionidae). *Iran. J. Ichthyol.*, 2021, 8(3):250-270.
11. Ghanbari M, Kneifel W, Domig KJ. A new view of the fish gut microbiome: Advances from nextgeneration sequencing. *Aquaculture*, 2019, 448:464-475.
12. Heidari Salkhordeh M R, Gharaei A, Mirdar Harijani J. First record of karyotype analysis in Anjak, *Schizocypris altidorsalis* (Bianco and Banareescu, 1982) from Hamoun Lake, Iran. *Iran. J. Fish. Sci.*, 2016, 15(1):246-254.
13. Heidari A, Sattari M, Mousavi-Sabet H, Alavi-Yeganeh MS. Morphological Comparison of two populations of Lake Goby *Rhinogobius similis* Gill, 1859 from Hariroud basin. *J. Appl. Ichthyol. Res.*, 2019, 7(1):143-154.
14. Loy A, Cataudella S, Corti M. Shape changes during the growth of the sea bass, *Dicentrarchus labrax* (Teleostea: Perciformes), in relation to different rearing conditions. In *Advances in Morphometrics*, Boston, MA: Springer US. 1996, pp 399-405.
15. Slice DE, Bookstein FL. A glossary for geometric morphometrics. Marcus LF, Corti M, Loy A, Naylor GJP, Slice D (Eds.), In *Advances in Morphometrics*, Boston, MA: Springer US. 1996, pp 531-551.
16. Mouludi-Saleh A, Eagderi S. Habitat-Associated Morphological Divergence of *Gasterosteus aculeatus* in the Southern Caspian Sea Basin. *Iran. J. Sci. Technol. Trans. Sci.*, 2021, 45:121-125.
17. Mouludi-Saleh A, Keivany Y. Morphological diversity in three species of Chubs (*Squalius* spp.) populations in Iranian Basins. *Nova Biologica Reperta*, 2018, 5(2):192-204.
18. Mouludi-Saleh A, Keivany Y. Morphometric analysis of *Squalius namak* Khaefi et al. 2016 in Khaznagh and Ghare-Chai rivers. *Sri Lanka J. Aquat. Sci.*, 2018, 23(2):173-178.
19. Nasri M, Eagderi S, Farahmand H, Segherloo IH. Body shape comparison of *Cyprinion macrostomum* (Heckel, 1843) and *Cyprinion watsoni* (Day, 1872) using geometric morphometric method. *Int. J. Aquat. Biol.*, 2013, 1(5):240-244.
20. Nasri M, Eagderi S, Keivany Y, Farahmand H, Dorafshan S, Nezhadheydari H. Morphological diversity of *Cyprinion* Heckel, 1843 species (Teleostei: Cyprinidae) in Iran. *Iran. J. Ichthyol.*, 2018, 5(2):96-108.
21. Nieceza AG. Morphological variation between geographically disjunct populations of Atlantic salmon: the effects of ontogeny and habitat shift. *Funct. Ecol.*, 1995, 9:448-456.

22. Pinheiro A, Teixeira CM, Rego AL, Marques JF, Cabral HN. Genetic and morphological variation of *Solea lascaris* (Risso, 1810) along the Portugal coast. *Fish. Res.*, 2005, 73:67-78.
23. Radkhah A, Poorbagher H, Eagderi S. Investigation of morphological differences of *Capoeta capoeta* populations in the upstream and downstream of Zarinerood River in Urmia Lake Basin. *J. Anim. Environ.*, 2016, 8(3):167-174.
24. Razavipour P, Eagderi S, Poorbagher H, Javanshir Khooi A, Keivany Y. Comparative study of morphological characteristics of Tuini fish (*Capoeta damascina*) in inland water of Iran using geometric morphometric method. *J. Fish.*, 2015, 8(1):79-90.
25. Rohlf FJ, Bookstein FL, (Eds). Proceedings of the Michigan Morphometrics Workshop. Museum of Zoology special publication no. 2, Ann Arbor: University of Michigan, 1990.
26. Rohlf FJ, Slice, DE. Extensions of the procrustes method for the optimal superimposition of landmarks. *Syst. Zool.*, 1990, 39(1):40-59.
27. Salehinia D, Eagderi S, Khorasani NA, Zamani-Faradonbeh, M. Impact of Sangban Dam on the morphological characteristics of Siah mahi (*Capoeta gracilis*, keyserling, 1864) populations using traditional and geometric morphometrics techniques. *J. Anim. Environ.*, 2016, 8(2):97-104.
28. Sansom RS. Phylogeny, classification, and character polarity of the Osteostraci (Vertebrata). *J. Syst. Palaeontol.*, 2009, 7:95-115.
29. Zamani Faradonbeh M, Eagderi S, Nasri M. Geometric morphometric comparison of populations of waspi *Cabdio morar* (Hamilton, 1822) in Mashkid and Mokran basins. *Iran. J. Fish. Sci.*, 2014, 23(2):57-67.

**How to cite this article:** Ganjali Z. Comparative Survey of Geometric Morphometrics and Traditional Methods in Male and Female of Anjak, *Schizocypris altidorsalis* (Bianco and Banareescu, 1982) from Sistan Basin, Iran. *Curr. Appl. Sci.*, 2025, 3(1):21-28.  
<https://doi.org/10.22034/cas.2023.410928.1032>